

Literatur zum Beitrag von Silber/Haude

- 1 Albiero R., Nishida T., Adamian M. et al.: Edge Restenosis After Implantation of High Activity ^{32}P Radioactive β -Emitting Stents. *Circulation*, 2000; 101: 2454-2457
- 2 Brenner D.J., Miller R.C.: Long-Term Efficacy of Intracoronary Irradiation in Inhibiting In-Stent Restenosis. *Circulation*, 2001; 103: 1330-1332
- 3 Buchbinder M., Strathearn G., Tarn L.A. et al.: RDX™ Radiation Delivery System: Balloon-Based Radiation Therapy. In: *Handbook of Vascular Brachytherapy, 2nd*, eds. R. Waksman and P. W. Serruys. London, Martin Dunitz Ltd., 1998; 145-152
- 4 Casterella P.J., Teirstein P.S.: Prevention of coronary restenosis. *Cardiol. Rev.*, 1999; 7: 19-31
- 5 Chae I.C., Kim H.S., Sohn D.W. et al.: Intracoronary Beta Radiation Therapy with ^{188}Re -DTPA-Filled Balloon System after Angioplasty - SPARE (Seoul National University Hospital Post-Angioplasty Rhenium) Trial. *Circulation*, 2000; 102: 11-751
- 6 Condado J.A., Lansky A.J., Saucedo J. et al.: Five Year Clinical and Angiographic Follow-Up after Intracoronary ^{192}Ir -Iridium Radiation Therapy. *Circulation*, 2000; 102: 11-750
- 7 Condado J.A., Waksman R., Gurdiel O. et al.: Long-term angiographic and clinical outcome after percutaneous transluminal coronary angioplasty and intracoronary radiation therapy in humans. *Circulation*, 1997; 96: 727-732
- 8 Costa M. A., Sabate M., van der Giessen W.J. et al.: Late Coronary Occlusion After Intracoronary Brachytherapy. *Circulation*, 1999; 100: 789-792
- 9 Coussement P., Stella P., Vanbilloen H. et al.: Intra-Coronary Radiation Therapy With Liquid Rhenium-186. *The RadioCathTrial*. *JACC*, 2000; 35: 1A
- 10 Dietz R., Silber S., Baumgart D. et al.: Positionspapier zur Intrakoronaren Brachytherapie. Stellungnahme der Deutschen Gesellschaft für Kardiologie- Herz und Kreislaufforschung, bearbeitet im Auftrag der Kommission für Klinische Kardiologie. *Z. Kardiol.*, 2001; 90: 875-880
- 11 Fischell T.A., Khanna B.K., Fischell D.R. et al.: Low-dose, beta-particle emission from „stent“ wire results in complete, localized inhibition of smooth muscle cell proliferation. *Circulation*, 1994; 90: 2956-2963
- 12 Giap H., Teirstein P., Massullo V. et al.: Barotrauma Due To Stent Deployment In Endovascular Brachytherapy for Restenosis Prevention. *Int. J. Radial. Oncol. Biol. Phys.*, 2000; 47: 1021-1024
- 13 Hall E.J.: DNA Strand breaks and chromosomal aberrations. In: Hall EJ ed. *Radiobiology for the Radiologist*. Philadelphia, Penn: J. B. Lippincott Company, 1994; 15-73
- 14 Hausleiter J., Li A., Knapp F.F.R. et al.: Low Body Exposure in Case of Radioactive Balloon Leakage - A Biodistribution and Elimination Study of ^{188}Re in Pigs. *JACC*, 1999; 33: 44A
- 15 Hehrlein C., Brachmann J., Hardt S. et al.: P-32 Stents for Prevention of Restenosis: Results of the Heidelberg Safety Trial Using the Palmaz-Schatz Stent Design at Moderate Activity Levels in Patients with Restenosis after PTCA. *Circulation*, 1998; 98: 1-780
- 16 Hehrlein C., De Vries J.J., Arab A. et al.: Failure of a Novel Balloon-Expandable γ -Emitting (^{103}Pd) Stent to Prevent Edge Effects. *Circulation*, 2001; 104: 2358-2362
- 17 Hehrlein C., Kollum M., Arab A. et al.: Increased Apoptotic Cell Death in the Neointima after Stent-Based Vascular Irradiation: Role of Radiation-Induced Apoptosis for Restenosis Reduction. *J. Interv. Cardiol.*, 1999; 12: 299-304
- 18 Hehrlein C., Kovacs A., Wolf G.K. et al.: A novel balloon angioplasty catheter impregnated with beta-particle emitting radioisotopes for vascular brachytherapy to prevent restenosis. First in vivo results. *Eur. Heart. J.*, 2000; 21: 2056-2062

- 19 Höher M., Kotzerke J., Wöhrle J. et al.: Intracoronary β -irradiation with liquid rhenium-188 to prevent restenosis following angioplasty of de-novo and restenotic lesions - results from the randomized ECRIS II trial. *Eur. Heart J.*, 2001; 22 (Suppl.): 5
- 20 Ikeno F., Carter A.J.: Will There Always Be an Edge to Radiation for the Prevention of Restenosis? *Cathet. Cardiovasc. Intervent.*, 2001; 54: 49-50
- 21 John M.C., Färb A., Virmani R.: Adverse Edge Effects Are Not Prevented by ^{32}P Beta-Emitting Hot-Ends Radioactive Stents. *JACC*, 2001; 37: 62 A
- 22 Kay I.P., Wardeh A.J., Kozuma K. et al.: The pattern of restenosis and vascular remodelling after cold-end radioactive stent Implantation. *Eur. Heart J.*, 2001; 22: 1311-1317
- 23 Kim H.S., Waksman R., Göttin Y. et al.: Edge Stenosis and Geographical Miss Following Intracoronary Gamma Radiation Therapy for In-Stent Restenosis. *J. Am. Coll. Cardiol.*, 2001; 37: 1026-1030
- 24 Kosuga K., Tamai H., Ueda K. et al.: Effectiveness of tranilast on restenosis after directional coronary atherectomy. *Am. Heart J.*, 1997; 134: 712-718
- 25 Kuntz R.E., Speiser B., Joyal M. et al.: Clinical and Angiographic Outcomes After Use of Sr-90 Beta Radiation for the Treatment of De Novo and Restenotic Coronary Lesions. Late breaking clinical trials, 73. Jahrestagung des AHA, New Orleans, LA, 2000
- 26 Leon M.B., Teirstein P.S., Moses J.W. et al.: Localized Intracoronary Gamma-Radiation Therapy To Inhibit The Recurrence of Restenosis After Stenting. *N. Engl. J. Med.*, 2001; 344: 250-256
- 27 Post M., Borst C., Kuntz R.E.: The relative importance of arterial remodeling compared with intimal hyperplasia in lumen renarrowing after balloon angioplasty: A study in the normal rabbit and the hypercholesterolemic Yucatan micropig. *Circulation*, 1994; 89: 2816-2821
- 28 Raizner A.E., Oesterle S.N., Waksman R. et al.: Inhibition of restenosis with extra-emitting radiotherapy; Report of the Proliferation Reduction with Vascular Energy Trail (PREVENT). *Circulation*, 2000; 102: 951-958
- 29 Reilly J.P., Grise M.A., Mehran R. et al.: SCRIPPS II: A Pilot Trial of Localized Radiation Therapy (Brachytherapy) to Inhibit Restenosis after Stenting. *Circulation*, 2001; 104: II-546
- 30 Rubin P., Soni A., Williams J.P.: The molecular and cellular biologic basis for the radiation treatment of benign proliferative diseases. *Semin. Radial. Oncol.*, 1999; 9: 203-214
- 31 Sabate M., Costa M.A., Kozuma K. et al.: Geographie Miss. A Cause of Treatment Failure in Radio-Oncology Applied to Intracoronary Radiation Therapy. *Circulation*, 2000; 101: 2467-2471
- 32 Schulz C., Niederer C., Andres M.S. et al.: Endovascular irradiation from b-particle-emitting gold Stents result in increased neointima formation in a porcine restenosis model. *Circulation*, 2000; 101: 1970-1975
- 33 Serruys P.W., Kay I. P.: I Like the Candy, I Hate the Wrapper. The ^{32}P Radioactive Stent. *Circulation*, 2000; 101: 3-7
- 34 Serruys P.W., Kutryk M.J.B., Bruining N. et al.: Antisense oligonucleotide against c-myc administered with the Transportos delivery catheter for the prevention of in-stent restenosis. Results of the randomized ITALICS trial. *Circulation*, 1998; 98: I-363
- 35 Sianos G., Kay I.P., Costa M.A. et al.: Geographical Miss During Catheter-Based Intracoronary Beta-Radiation: Incidence and Implications in the BRIE Study. *J. Am. Coll. Cardiol.*, 2001; 38: 415-420
- 36 Silber S.: Für die START-Studienteilnehmer. *Die START-Studie. Z. Kardiol.*, 2000; 89: 185
- 37 Silber S.: Antiproliferativ beschichtete Stents und Brachytherapie: Gemeinsamkeiten und Unterschiede. *Z. Kardiol.*, 2002 (im Druck)
- 38 Silber S., Brockhoff C., Doerr R. et al.: Notwendigkeit für die einjährige Gabe von Clopidogrel zur Vermeidung später Stentthrombosen nach in-trakoronarer Brachytherapie: erste Ergebnisse des Deutschen IST-Reg-isters. *Z. Kardiol.*, 2001; 90: 34

- 39 Silber S., Krischke I., Tourlakidou S. et al.: Kombinierte Anwendung von Brachytherapie und Cutting-Balloon zur Therapie rezidivierender In-Stent-Restenosen. *Z. Kardiol.*, 2001; 90: 313
- 40 Silber S., Serruys P.W., Bonnier J. et al.: Die Lernkurve bei intrakoronarer Brachytherapie und ihr Einfluss auf das angiographische 6-Monats-Ergebnis: endgültige Ergebnisse der Europäischen BRIE (Beta Radiation In Europe)-Studie. *Z. Kardiol.*, 2001; 90 (Suppl. 2): 219
- 41 Silber S., von Rottkay P., Gielow A. et al.: Intrakoronare Brachytherapie mit Strontium/Yttrium-90. *Herz*, 1998; 23: 380-393
- 42 Silber S., von Rottkay P., Lössl L. et al.: Sicherheit und Durchführbarkeit der intrakoronaren Brachytherapie mit dem Novoste™-System im Rahmen von internationalen Multicenter-Studien. *Z. Kardiol.*, 2000; 89: 323-329
- 43 Teirstein P.S., Kuntz R.E.: New Frontiers in Interventional Cardiology. In-travascular Radiation to Prevent Restenosis. *Circulation*, 2001; 104: 2620-2626
- 44 Teirstein P.S., Massullo V., Jani S. et al.: Catheter-based radiotherapy to inhibit restenosis after coronary Stenting. *N. Engl. J. Med.*, 1997; 336: 1697-1703
- 45 Teirstein P.S., Massullo V., Jani S. et al.: Three-Year Clinical and Angiographic Follow-up After Intracoronary radiotherapy. Results of a Randomized Clinical Trial. *Circulation*, 2000; 101: 360-365
- 46 Teirstein P.S., Massullo V., Jani S. et al.: Two-Year Follow-Up After Catheter-Based Radiotherapy to Inhibit Coronary Restenosis. *Circulation*, 1999; 99: 243-247
- 47 Tripuraneni P., Parikh S., Giap H. et al.: How Long Is Enough? Defining the Treatment Length in Endovascular Brachytherapy. *Cathet. Cardiovasc. Intervent*, 2000; 51: 147-153
- 48 Van der Giessen W.J., Regar E., Hartevelde M.S. et al.: „Edge Effect“ of ³²P Radioactive Stents Is Caused by the Combination of Chronic Stent Injury and Radioactive Dose Falloff. *Circulation*, 2001; 104: 2236-2241
- 49 Von Dahl J., Haager P.K., Reineke T. et al.: Predictors of Recurrent In-Stent Restenosis Following Mechanical Treatment by Angioplasty or Rotational Atherectomy: Results From an Angiographically Controlled Prospective Trial (ARTIST Study). *J. Am. Coll. Cardiol.*, 2000; 35: 83A
- 50 Waksman R., Ajani A.E., Kim H.S. et al.: Twelve versus Six Months of Clopidogrel to Prevent Late Total Occlusion after Gamma Radiation Therapy for In-Stent Restenosis: WRIST 12 vs. WRIST PLUS. *Circulation*, 2001; 104: 11-577
- 51 Waksman R., Ajani A.E., White L.R. et al.: Two-Year Follow-Up After Beta-and Gamma Intracoronary Radiation Therapy for Patients With Diffuse In-Stent Restenosis. *Am. J. Cardiol.*, 2001; 88: 425-28
- 52 Waksman R., Ajani A.E., White R.L. et al.: Prolonged Antiplatelet Therapy to Prevent Late Thrombosis After Intracoronary γ -Radiation in Patients With In-Stent Restenosis. Washington Radiation for In-Stent Restenosis Trial Plus 6 Months of Clopidogrel (WRIST PLUS). *Circulation*, 2001; 103: 2332-2335
- 53 Waksman R., Bhargava B., Mintz G.S. et al.: Late Total Occlusion After Intracoronary Brachytherapy for Patients With In-Stent Restenosis. *J. Am. Coll. Cardiol.*, 2000; 36: 65-68
- 54 Waksman R., Elsayyad S., Mehran R. et al.: High Dose Intracoronary Gamma Radiation for Patients with Diffuse In-Stent Restenosis. *Circulation*, 2000; 102: 11-667
- 55 Waksman R., Raizner A., Lansky A.J. et al.: Beta Radiation to Inhibit Recurrence of In-Stent Restenosis: Study Design, Device and Dosimetry Details of the Multicenter Randomized Double Blind Study. *Circulation*, 2000; 102: 11-667
- 56 Waksman R., Robinson K.A., Crocker I.R. et al.: Endovascular Low-Dose Irradiation Inhibits Neointima Formation After Coronary Artery Balloon In-jury in Swine. A Possible Role for Radiation Therapy in Restenosis Prevention. *Circulation*, 1995; 91: 1533-1539
- 57 Waksman R., Rodriguez J.C., Robinson K.A. et al.: Effect of Intravascular Irradiation on Cell Proliferation, Apoptosis, and Vascular Remodeling After Balloon Overstretch Injury of Porcine Coronary Arteries. *Circulation*, 1997; 96: 1944-1952
- 58 Waksman R.: The SVG-WRIST trial, Symposiumsreferat, 50. Jahrestagung des ACC. Orlando, FL, 2001

- 59 Waksman R.: The LONG-WRIST trial, Symposiumsreferat, 49. Jahrestagung des ACC. Anaheim, CA, 2000
- 60 Waksman R., White L., Chan R.C. et al.: Intracoronary γ -Radiation Therapy After Angioplasty Inhibits Recurrence in Patients With In-Stent Restenosis. *Circulation*, 2000; 101: 2165-2171
- 61 Waksman R.: Late Thrombosis After Radiation - Sitting on a Time Bomb. *Circulation*, 1999; 100: 780-782
- 62 Waksman R.: Response to radiation therapy in animal restenosis models. *Semin. Intervent. Cardiol.*, 1997; 2: 95-101
- 63 Weinberger J., Amols H., Ennis R.D. et al.: Intracoronary Irradiation: Dose response for the prevention of restenosis in swine. *Int. J. Radiat. Oncol. Biol. Phys.*, 1996; 36: 767-775
- 64 Wiedermann J.G., Marboe C., Amols H. et al.: Intracoronary Irradiation markedly reduces restenosis after balloon angioplasty in a porcine model. *J. Am. Coll. Cardiol.*, 1994; 23: 1491-1498
- 65 Wiedermann J.G., Marboe C., Amols H. et al.: Intracoronary Irradiation Makedly Reduces Neointimal Proliferation After Balloon Angioplasty in Swine: Persistent Benefit at 6-Month Follow-Up. *J. Am. Coll. Cardiol.*, 1995; 25: 1451-1456